






SYSTEMATIC REVIEW

Prevalence of periodontitis in dentate people between 2011 and 2020: A systematic review and meta-analysis of epidemiological studies

Diogo Trindade¹ | Rui Carvalho¹  | Vanessa Machado^{1,2}  |
Leandro Chambrone^{2,3,4}  | José João Mendes¹  | João Botelho^{1,2} 

¹Clinical Research Unit (CRU), Egas Moniz Center for Interdisciplinary Research, Egas Moniz School of Health & Science, Almada, Portugal

²Evidence-Based Hub, Egas Moniz Center for Interdisciplinary Research, Egas Moniz School of Health & Science, Almada, Portugal

³Unit of Basic Oral Investigation (UIBO), Universidad El Bosque, Bogotá, Colombia

⁴Department of Periodontics, School of Dental Medicine, The University of Pennsylvania, Philadelphia, Pennsylvania, USA

Correspondence

João Botelho, Quinta da Granja, Campus Universitário, Monte da Caparica, 2829-511, Almada, Portugal.
Email: jbotelho@egasmoniz.edu.pt

Abstract

Aim: The aim of the study was to evaluate the prevalence of periodontitis in dentate people between 2011 and 2020.

Materials and Methods: PUBMED, Web of Science, and LILACS were searched up to and including December 2021. Epidemiological studies reporting the prevalence of periodontitis conducted between 2011 and 2020 were eligible for inclusion in this review. Studies were grouped according to the case definition of confidence as confident (Centers for Disease Control [CDC] AAP 2012; CDC/AAP 2007; and Armitage 1999) and non-confident (community periodontal index of 3 or 4, periodontal pocket depth >4 mm, and clinical attachment level ≥ 1 mm). Random effects meta-analyses with double arcsine transformation were conducted. Sensitivity subgroup and meta-regression analyses explored the effect of confounding variables on the overall estimates.

Results: A total 55 studies were included. The results showed a significant difference, with confident case definitions (61.6%) reporting nearly twice the prevalence as non-confident classifications (38.5%). Estimates using confident periodontal case definitions showed a pooled prevalence of periodontitis of 61.6%, comprising 17 different countries. Estimates reporting using the CDC/AAP 2012 case definition presented the highest estimate (68.1%) and the CDC/AAP 2007 presented the lowest (48.8%). Age was a relevant confounding variable, as older participants (≥ 65 years) had the highest pooled estimate (79.3%).

Conclusion: Between 2011 and 2020, periodontitis in dentate adults was estimated to be around 62% and severe periodontitis 23.6%. These results show an unusually high prevalence of periodontitis compared to the previous estimates from 1990 to 2010.

KEYWORDS

periodontal disease, periodontitis, prevalence, systematic review

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Journal of Clinical Periodontology* published by John Wiley & Sons Ltd.

Clinical Relevance

Scientific rationale for study: To comprehensively estimate the prevalence of periodontitis reported in studies between 2011 and 2020 sourced from epidemiological data.

Principal findings: During this period, periodontitis was collectively estimated to be around 63% and severe periodontitis 24.2%. Pooled estimates from confident case definitions presented almost two-fold prevalence.

Practical implications: These results highlight the need for strengthening the surveillance of periodontitis and informing global health stakeholders to devise and better manage public periodontal health strategies.

1 | INTRODUCTION

Periodontitis is a public health problem whose high prevalence contributes to the global burden of chronic non-communicable diseases (Petersen, 2009; Tonetti et al., 2017; Caton et al., 2018; Tonetti et al., 2018). According to the Global Burden of Disease (GBD), periodontitis was ranked as one of the most prevalent conditions of humankind between 1990 to 2010 (Marcenes et al., 2013; Kassebaum et al., 2014), and a recent update until 2019 confirmed that this prevalence is still substantial and worrisome (Wu et al., 2022).

Periodontitis had a significant socio-economic impact in the United States and Europe in 2018 of around \$154 billion and €159 billion, respectively (Botelho et al., 2022). Also, its impact on patients' quality of life and systemic health has been extensively verified over the last decades (Buset et al., 2016; Botelho et al., 2020, 2022).

The prevalence of periodontitis has been reported using different and highly heterogeneous approaches. On one hand, prevalence data provided by the GBD resort to arithmetic extrapolations based on miscellaneous periodontal classifications (Schwendicke et al., 2018; GBD 2017 Oral Disorders Collaborators et al., 2020; Botelho et al., 2022; Wei et al., 2021; Wu et al., 2022), whose reliability is under debate due to the variety of case definitions (Marcenes et al., 2013; Kassebaum et al., 2017; GBD 2017 Oral Disorders Collaborators et al., 2020). On the other hand, methodological heterogeneity hampers a traceable comparison over time (Frencken et al., 2017). Thus, it is imperative to carry out continuous studies of estimates, such as summary measures of population health (e.g., disability-adjusted life years), or through meta-analysis.

Considering the challenges with current measurement methodologies for periodontitis, measurable specific estimates based on epidemiological studies are warranted to advance global public health (Kassebaum et al., 2017). Therefore, this systematic review aims to comprehensively estimate the prevalence of periodontitis reported in studies between 2011 and 2020 sourced from epidemiological data. Our secondary objectives are (i) to evaluate the prevalence geographically, (ii) to compare estimates from confident and non-confident case definitions, and (iii) to explore other confounding variables. The following focused PECO question was addressed: "What is the pooled prevalence estimate of periodontitis in epidemiological studies carried out between 2011 and 2020?" (Population: among dentate adults assessed in an epidemiological survey; Exposure: periodontitis; Comparison: periodontal status assessed; Outcome: prevalence).

2 | METHODS

This systematic review protocol was previously set by all the authors and was registered in the National Institute for Health Research PROSPERO, the International Prospective Register of Systematic Reviews (<http://www.crd.york.ac.uk/PROSPERO>, ID Number: CRD42021231357). The review design followed the Cochrane Handbook of Systematic Reviews of Interventions (Higgins et al., 2019) and was reported according to the updated Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (M. J. Page et al., 2021). The PRISMA checklist is provided as Supplementary File 1.

2.1 | Type of studies, type of intervention, and inclusion criteria

The eligibility criteria were as follows: observational epidemiological studies (cross-sectional and cohort studies) reporting the prevalence of periodontitis in adults (18 years old or older) between 2011 and 2020. Cross-sectional and cohort studies (data were gathered from baseline reports) represent epidemiological studies in nature and provided prevalence traits important for the aims of this systematic review. Case-control studies were not considered because of the associated inconvenience in sampling cases and controls (Belbasis & Bellou, 2018). In line with this, intervention studies (both randomized and non-randomized) were considered only if they had a baseline inclusion of participants with an epidemiological approach.

In contrast, studies reporting the prevalence of periodontitis of a specific population (e.g., pregnant women, people with a particular underlying disease, etc.), studies based on self-reported case definitions of periodontitis, and pre-clinical studies were excluded.

In cases where additional clarifications were required, the corresponding author of the included study was contacted via e-mail, and, in case of no response, again 1 week later.

2.2 | Primary and secondary outcomes

The primary outcome of this systematic review was the prevalence of periodontitis reported in epidemiological studies, either national or regional. We further explored the distribution of periodontitis per

continent, based on the case definition, according to established age intervals, as well as the severity of periodontitis.

2.3 | Information sources search

The search strategy sought to identify all studies reporting the prevalence of periodontitis conducted between January 2011 and December 2020. We defined this year interval, based on the recent estimates by Kassebaum et al. (2017) for the period between 1990 and 2010.

Detailed search strategies were used, without language restrictions, on the following electronic databases: PubMed, Web of Science, and LILACS (Latin-American Scientific Literature in Health Sciences). The search algorithm, developed using keywords and the Medical Subject Headings (MeSHs), was as follows: “(periodont* OR ‘chronic periodontitis’ OR (periodontal diseases [MeSH]) OR ‘attachment loss’ OR pocket*) AND (prevalence [MeSH] OR epidemiology [MeSH])”.

For the remaining databases, the search was adapted accordingly.

2.4 | Study selection

Two researchers (D.T. and R.C.) independently selected the relevant articles by screening the titles and abstracts, excluding the non-relevant studies. Any paper classified as potentially eligible by either reviewer was appraised by a full-text reading and the reasons for exclusion were fully detailed. Any disagreement was resolved through discussion with a third reviewer (J.B.) and a decision arrived at by consensus.

2.5 | Data extraction process and data items

The studies that fulfilled the inclusion criteria were organized into evidence tables describing the characteristics and results of each study, including the following: study identification (i.e., first author's name and publication year), time period of the study, continent, country of origin of the research, country coordinates, funding information, inclusion and exclusion criteria, periodontal case definition, characteristics and number of participants, and outcome measures. We later added the methodological risk of bias of the study (detailed in Section 2.6). All disagreements were resolved through discussion with a third reviewer (J.B.). Because of the known impact of case definitions in prevalence estimates (Holtfreter et al., 2015), we categorized case definitions following the strategy used by Muñoz Aguilera et al. (2020) as confident and non-confident, as explained below.

2.5.1 | Confident case definition of periodontitis

The following case definitions were considered as confident:

- Interdental clinical attachment loss (CAL) in ≥ 2 non-adjacent teeth, or buccal or oral CAL ≥ 3 mm with periodontal pocket depth (PPD)

> 3 mm detectable at ≥ 2 teeth (American Academy of Periodontology [AAP]/European Federation of Periodontology [EFP]; Tonetti et al., 2018);

- Two or more inter-proximal sites with CAL ≥ 3 mm and two or more inter-proximal sites with PPD ≥ 4 mm (not on the same tooth) or one site with PPD ≥ 5 mm (Centers for Diseases Control [CDC]/AAP, 2012; Eke et al., 2012);
- At least two sites on different teeth with clinical attachment level (CAL) 6 mm and at least one site with PPD 4 mm (CDC/AAP 2007; R. C. Page & Eke, 2007);
- Generalized chronic periodontitis (at least 30% sites with CAL ≥ 4 mm; CDC 1999; Armitage, 1999).

2.5.2 | Non-confident case definition of periodontitis

The following reported criteria were considered as a non-confident case definition: the (WHO) CPI score 3/4 in at least one quadrant; at least one site with PPD > 4 mm; CAL ≥ 1 mm.

2.6 | Risk-of-bias assessment

The methodological quality of the included studies was carried out by two independent reviewers (D.T. and R.C.) using the “Assessing risk of bias in population-based prevalence studies” tool (Hoy et al., 2012). Any disagreement was resolved through discussion with a third reviewer (J.B.).

The tool consists of 10 items addressing external validity (items 1–4, accounting for the selection and non-response bias domains) and internal validity (items 5–10, accounting for the measurement bias and bias related to the analysis domains). Each item was rated as having either “yes” (low) or “no” (high) risk of bias. Items with insufficient information to properly meet the requirements were classified as “no” (i.e., high risk of bias; Hoy et al., 2012). Following Hoy et al., the overall risk of bias of each study was decided on the basis of agreement between all authors. Therefore, each article's overall risk of bias was evaluated as “High” (i.e., if only 0–3 items were responded with yes, having an important impact on our confidence in the estimate, and will likely change the estimate), “Moderate” (if 4–8 items were responded with yes), and “Low” (if 9 or more items were responded with yes, indicating that further research is very unlikely to change our confidence in the estimate).

2.7 | Statistical analysis

Extracted data were organized into evidence tables. Because of the existence of multiple categories of prevalence for periodontitis in the literature, and the possible imbalance in weight of the studies affecting the meta-analysis, we implemented a double arcsine transformation meta-analysis (Barendregt et al., 2013). Considering the




FIGURE 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses 2020 flow diagram for new systematic reviews that included searches of databases and registers only. Consider, if feasible to do so, reporting the number of records identified from each database or register searched (rather than the total number across all databases/register). If automation tools were used, indicate how many records were excluded by a human and how many were excluded by automation tools. Source: M. J. Page et al. (2021). For more information, visit: <http://www.prisma-statement.org/>. [Colour figure can be viewed at wileyonlinelibrary.com]

geographical variation and difference in populations, we could not assume the existence of a true effect size; so we employed a random-effects model (Schwarzer et al., 2015), as previously described (Schwarzer, 2007). All random-effects meta-analysis and forest plots were performed in R version 3.4.1 using the *meta* package (Schwarzer, 2007). The results are presented as percentage prevalence ($p \times 100\%$) and 95% confidence intervals (CI). Heterogeneity was explored through the I^2 index and Cochrane's Q statistic ($p < .1$) and χ^2 test for the overall homogeneity (Higgins et al., 2019). Substantial heterogeneity was defined as $I^2 > 50\%$. All tests were two-tailed, with α set at .05.

We conducted a series of a priori sensitivity analyses (i.e., subgroup analyses) to examine the effect of the case definition of confidence (confident vs. non-confident) and risk of bias (low vs. moderate-to-high) to the overall estimates. Meta-regression was used to explore adjusting effects, particularly confounding variables,

on the prevalence estimates, particularly female/male ratio, latitude, longitude, and study sample size.

3 | RESULTS

3.1 | Study selection

The electronic searches retrieved a total of 5756 records (1943 from PubMed, 3023 from Web of Science, and 790 from LILACS; Figure 1). After removal of duplicates, 4968 articles were excluded based on title/abstract assessment, and 28 articles were unable to be accessed. Out of the articles remaining, 379 were excluded in the full-text appraisal, due to not meeting the inclusion criteria (with the respective reason detailed in Supplementary File 2). Finally, a total of 57 articles (Moya et al., 2012; Figueiredo et al., 2013; Araya Vallespir

et al., 2014; Jaafar et al., 2014; Juarez et al., 2014; Thanakun et al., 2014; Bhat et al., 2015; Eke et al., 2015; Giacaman et al., 2016; Khan et al., 2016; Holde et al., 2017; Ramírez et al., 2017; Shyagali et al., 2017; Silva-Junior et al., 2017; Zaitso et al., 2017; Balaji et al., 2018; Bhat et al., 2018; Eke et al., 2018; He et al., 2018; Iwasaki et al., 2018; Ortiz et al., 2018; Pinto-Filho et al., 2018; Shariff et al., 2018; Skośkiewicz-Malinowska et al., 2018; Wahlin et al., 2018; Botelho et al., 2019; Dhaifullah et al., 2019; Helmi et al., 2019; Lasta et al., 2019; J. B. Lee et al., 2019; K. Lee & Kim, 2019; Shimizu et al., 2019; Zhao et al., 2019; Bongo et al., 2020; Díaz-Reissner et al., 2020; Ha et al., 2020; Nakamura et al., 2020; Romero-Castro et al., 2020; Schmidt et al., 2020; Sekiguchi et al., 2020; Singh et al., 2020; Sun et al., 2020; Bilgin Çetin et al., 2021; Clauss et al., 2021; Costa et al., 2021; Germen et al., 2021; Ghassib et al., 2021; Goel et al., 2021; Gomes-Filho et al., 2021; Han & Kim, 2021; Iwasaki et al., 2021; Jiao et al., 2021; Kocher et al., 2021; Oliveira et al., 2021; Sakurai et al., 2021; Song et al., 2021; Stødle et al., 2021; Su et al., 2021) were included in the qualitative and quantitative analyses of the present systematic review. Two studies had their data reported in more than one article, so these papers were grouped under a single name (Bhat et al., 2015; Eke et al., 2015; Bhat et al., 2018; Eke et al., 2018). Therefore, 55 articles were finally included. Inter-examiner reliability at full-text screening was considered excellent (κ [kappa] score = 0.86, 95% CI: 0.83–0.89).

One study did not report data for the overall prevalence of periodontitis, but it reported data for the subgroup analysis information of moderate to severe periodontitis (Ha et al., 2020). To complete information on the year range of interest, we used the official report where this information was detailed (Australian Research Centre for Population Oral Health, 2019).

3.2 | Study characteristics

Studies from 25 different countries across Asia, America, Europe, Australia, and Africa were included (Table 1). Most studies ($n = 30$, 54.5%) reported periodontal outcomes using a non-confident case definition, either the community periodontal index (CPI) score 3/4 in at least one quadrant, at least one site with PPD >4 mm, or CAL \geq 1 mm. Within the confident periodontal definitions, the CDC/AAP 2012 was the most employed, being reported in 15 studies (27.2%); the remaining reported periodontitis based on confident case definitions.

3.3 | Risk of bias

Forty-four studies were considered of moderate risk of bias, while only 11 were of low risk of bias (Supplementary File 3). Overall, studies failed to report prevalence data that represented closely the national population regarding relevant variables (81.8%, $n = 45$) and to use an acceptable case definition of periodontitis (45.5%, $n = 25$). Non-response bias was minimal (78.2%, $n = 43$), while 61.7% ($n = 50$) used some form of random selection and 67.3% ($n = 37$) studies had

the prevalence period as the parameter of interest. The remaining items had predominantly low risk of bias (over 90% of the studies). Inter-examiner reliability at risk of bias assessment was considered excellent (κ score = 0.82, 95% CI: 0.79–0.84).

3.4 | Pooled estimates

Between 2011 and 2020, a total of 88,917 adults were included in the overall pool of analyses, with 44,614 adults reported to have periodontitis. We started by comparing non-confident with confident case definitions of periodontitis. Overall, the results confirmed a significant difference ($p < .00001$), with confident case definitions (61.6%, 95% CI: 55.1–67.9, $p < .000001$, $I^2 = 99.1\%$) reporting nearly twice the prevalence than non-confident classifications (38.5%, 95% CI: 30.4–46.9, $I^2 = 99.7\%$). For this reason, and following the instruction of GRADE, we focused on the estimates using confident periodontal case definitions (Schünemann et al., 2008).

Furthermore, we explored the impact of partial-mouth protocols and non-reported protocols in the overall estimates. Sensitivity analyses showed that studies using a full-mouth protocol reported nearly twice the prevalence (45.7%, 95% CI: 38.0–53.6, $p < .000001$, $I^2 = 99.4\%$, number of studies = 32) as partial-mouth protocols (55.1%, 95% CI: 44.3–65.6, $p < .000001$, $I^2 = 99.7\%$, number of studies = 19) or non-reported protocols (40.9%, 95% CI: 8.9–78.2, $p < .000001$, $I^2 = 99.8\%$, number of studies = 3). The comparison for subgroup differences was not significant ($p = .3528$).

3.4.1 | Total prevalence

The overall prevalence of periodontitis was estimated at 61.6% (95% CI: 55.1–67.9, $p < .000001$, $I^2 = 99.1\%$), comprising 17 different countries. This estimate was confirmed to be affected by the methodological quality ($p = .0497$), comparing studies of low risk of bias (54.1%, 95% CI: 47.4–60.6, $p < .000001$, $I^2 = 98.9\%$) and studies with moderate risk of bias (64.7%, 95% CI: 56.3–72.6, $p < .000001$, $I^2 = 99.0\%$). There was substantial heterogeneity.

The latitude (estimate = 0.000, SE = 0.002, $p = .805$), longitude (estimate = 0.000, SE = 0.001, $p = .774$), and sample size (estimate = 0.000, SE = 0.000, $p = .9410$) were confirmed to have no influence on the final estimates through meta-regression. In addition, no publication bias was detected (Egger test = 0.61, SE = 3.63, $p = .8671$).

Considering the case definition, there were significant differences between all four reported classifications ($p = .0101$; Table 2). Estimates using CDC/AAP 2012 case definition presented the highest estimate (68.1%), while those using the CDC/AAP 2007 presented the lowest (48.8%).

We further analysed whether the age interval of participants in the included studies would influence the results. Overall, the test for subgroup differences could not confirm a difference between the age

TABLE 1 Study characteristics

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response rate/ eligibility % (n) | Edentulous participants (n) | Periodontal criteria (confidence) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|---------------------------------------|--|---|--|----------------------------------|-----------------------------|-----------------------------------|----------------------------------|----------------------------|----------------------|--------------|----------------|
| Moya et al. (2012) (Chile) | ≥60 years old, beneficiary of the public health system, and assisted at the health center at East Santiago of Chile | NR | NR | NR | 7.2% | CPI (non-confident) | Partial-mouth (CPI) | 168/212 | ≥60 | NR | NR |
| Figueiredo et al. (2013) (Brazil) | Kiriri Indians aged 19 years and older who were living in an isolated Indian area in the state of Bahia, in Northeast Brazil | Cardiovascular diseases and other conditions that require antibiotics before periodontal probing | 1025 | 22.0 (225) | 0.0% | CDC/AAP 2007 (confident) | Full-mouth (six sites) | 85/130 | ≥19 | 96/119 | Research grant |
| Thanakun et al. (2014) (Thailand) | NR | Patients with systemic diseases, as well as those who had received medications | 125 | 100.0 (125) | 0.0% | Armitage 1999 (confident) | Full-mouth (six sites) | 90/35 | 35–76 | 53/72 | Research grant |
| Jaafar et al. (2014) (Malaysia) | ≥19 years old | NR | 586 | 100.0 (586) | 0.0% | CPI (non-confident) | Partial-mouth (CPI) | 228/309 | ≥19 | 218/319 | Research grant |
| Araya Vallespir et al. (2014) (Chile) | 35–44 years old, belonging to the family health center Lorenzo Arenas from the municipality of Concepción, Chile | NR | 58 | 100.0 (58) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (six sites) | 32/26 | 35–44 | NR | NR |
| Juarez et al. (2014) (Chile) | 18–70 years old, ≥2 teeth in the mouth | Pregnant, underwent periodontal treatment during the last 6 months, undergoing current periodontal treatment, on antibiotic treatment for a week or more in the last 6 months, or on treatment with immunosuppressive | 136 | 100.0 (136) | 0.0% | PPD ≥4 mm (non-confident) | Full-mouth (6 sites) | 67/69 | 18–70 | 51/85 | NR |

TABLE 1 (Continued)

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response rate/ eligibility % (n) | Edentulous participants (n) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|----------------------------------|---|---|---|---|-----------------------------------|--|-------------------------------|-------------------------|-----------------|-------------------|
| Khan et al. (2016) (Pakistan) | ≥18 years old | Participants who had received periodontal treatment in the last 4 months, participants on antibiotics within the past 4 months, pregnant women and lactating mothers, mentally handicapped, or on prophylactic antibiotics, or systemic/topical steroidal anti-inflammatory drugs for the last 4 months | 443 | 100.0 (443) | 0.0% | Partial-mouth (sextants) | 310/133 | ≥18 | 287/156 | Self-funded |
| Giacaman et al. (2016) (Chile) | The entire population of the Maule Region, divided according to the ages of epidemiological surveillance indicated by the WHO, that is, 6, 12, 15, 35–44, and 65–74 years | Systemic conditions that contraindicate periodontal evaluation, people who suffered heart disease, bleeding disorders or were under anticoagulant drug therapy, and all participants who were within the study ages but | 891 | 100.0 (891) | NR | Partial-mouth (CPI) | 162/531 | 35–44; 65–74 | 124/ 569 | Research grant |

(Continues)

TABLE 1 (Continued)

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response rate/ eligibility % (n) | Edentulous participants (n) | Periodontal criteria (confidence) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|-------------------------------------|---|---|--|----------------------------------|-----------------------------|-----------------------------------|----------------------------------|----------------------------|----------------------|--------------|----------------|
| Ramírez et al. (2017) (Colombia) | ≥35 years old, full clinical history and periodontal chart, presence of at least 10 teeth in the mouth excluding third molars | NR suffered from some cognitive deficit or mental disability | NR | NR | 0.0% | Armitage 1999 (confident) | Full-mouth (six sites) | 314/153 | ≥35 | 112/355 | NR |
| Zaitou et al. (2017) (Japan) | Adults aged 19–70 years old, employed at 11 companies (Company A–K) in the Kanto region of Japan | NR | 1078 | 100.0 (1078) | 0.0% | CPI (non-confident) | Partial-mouth (sextants) | 109/969 | 19–70 | 798/270 | Research grant |
| Silva-Junior et al. (2017) (Brazil) | Adults aged between 20 and 64 years | NR | 248 | 100.0 (248) | 0.0% | CPI (non-confident) | Partial-mouth (sextants) | 107/141 | 20–64 | NR | Research grant |
| Shyagali et al. (2017) (India) | Subjects who indulged in the use of tobacco products more than four times a week and for not less than a year | Subjects who consumed any of the tobacco products less than four times a week, who had a systemic disease, and who were on medication for such diseases | NR | NR | 0.0% | CPI (non-confident) | Partial-mouth (CPI) | 143/337 | 18–50 | NR | NR |
| Holde et al. (2017) (Norway) | Adults aged 20–79 years, living in Troms County, Norway | Edentulous or only one tooth | 2909 | 65.7 (1911) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (six sites) | 946/965 | 20–79 | 1016/895 | Research grant |
| He et al. (2018) (China) | Subjects within the 35–44, 55–64, and 65–74 years age groups, had the ability to understand and answer the | Subjects from 45 to 54 years age group, undergoing orthodontic treatment, teeth were covered by lots of calculus | 540 | 88.9 (480) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth | 296/184 | 35–44; 55–64; 65–74 | 243/237 | Research grant |

TABLE 1 (Continued)

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response participants/ rate/ eligibility % (n) | Edentulous participants (n) | Periodontal criteria (confidence) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|-----------------------------------|--|---|--|--|-----------------------------|-----------------------------------|----------------------------------|----------------------------|----------------------|--------------|----------------|
| Iwasaki et al. (2018) (Japan) | questionnaire, and had lived in the selected communities or villages for more than 6 months in the past year | Participants who had chronic hepatitis C infection and chronic hepatitis B infection | 1280 | 95.8 (1226) | 0.0% | PPD \geq 4 mm (non-confident) | Full-mouth (six sites) | 941/285 | 40–59 | NR | Self-funded |
| Ortiz et al. (2018) (Puerto Rico) | NR | History of clinically diagnosed diabetes, less than four natural teeth, a history of conditions that increase the risk of systemic complications during a periodontal exam, and inability to complete study procedures, fasting, two-hour glucose, or HbA1c levels met the American Diabetes Association thresholds for diabetes at the baseline exam | 773 | 95.1 (735) | 0.0% | CDC/AAP 2007 | Full-mouth (six sites) | 438/297 | 41–70 | 204/531 | Research grant |
| Bhat et al. (2015, 2018) (India) | Participants aged 35–54 years, dentate | Any participant reporting medical conditions (uncontrolled diabetes, heart disease, bleeding disorders) that contra-indicated periodontal probing | 1401 | 62.0 (869) | 0.0% | CDC/AAP 2012 | Full-mouth (six sites) | 405/464 | 35–54 | 472/397 | NR |

(Continues)

TABLE 1 (Continued)

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response eligibility % (n) | Edentulous participants (n) | Periodontal criteria (confidence) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|---|---|---|--|----------------------------|-----------------------------|---|----------------------------------|----------------------------|----------------------|--------------|----------------|
| Balaji et al. (2018) (India) | ≥18 years old | Participants who were pregnant, mentally ill, edentulous, non-ambulatory, or critically ill | 1000 | 100.0 (1000) | 0.0% | CAL ≥1 mm (non-confident) | NR | 423/577 | ≥18 | NR | Research Grant |
| Eke et al., (2015, 2018) (USA) | Adults 30 years or older who had one or more natural teeth and no health conditions requiring antibiotic prophylaxis before periodontal probing | Medical conditions, incomplete oral examinations, and institutionalized participants | 19,931 | 34.8 (6940) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (6 sites) | 3320/3620 | ≥30 | NR | NR |
| Pinto-Filho et al. (2018) (Brazil) | Kiriri Indians aged ≥19 years living in an isolated area in Bahia state, northeast Brazil | Patients with missing information | 226 | 99.6 (225) | 0.0% | Severe periodontitis ≥2 proximal sites CAL ≥6 mm, not on the same tooth, and ≥1 proximal site with a PPD ≥5 mm. (non-confident) | Full-mouth (six sites) | 65/160 | ≥19 | 101/124 | Research grant |
| Shariff et al. (2018) (USA) | ≥65 years old | NR | 964 | 89.2 (860) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (6 sites) | 691/169 | ≥65 | NR | Research grant |
| Skośkiewicz-Malinowska et al. (2018) (Poland) | ≥65 years old, local resident, able to communicate, and a written consent to participate in the survey | Coexisting systemic diseases in which dental pocket probing leading to transient bacteremia might have posed a risk for the patient's overall health condition: cardiovascular diseases (patients with heart valves, after heart transplant, with | 500 | 100.0 (500) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (6 sites) | 448/52 | ≥65 | NR | Research grant |

TABLE 1 (Continued)

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response participants/ rate/ eligibility % (n) | Edentulous participants (n) | Periodontal criteria (confidence) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|---|---|--|--|--|-----------------------------|-----------------------------------|--|----------------------------|----------------------|--------------|----------------|
| Wahlin et al. (2018) (Sweden) | 20–80 years old | congenital heart diseases, or with infective endocarditis), blood diseases (thrombocytopenia, haemophilia, von Willebrand disease), viral diseases (B and C type hepatitis, AIDS/HIV), as well as patients with Multi-Drug Resistant Organisms (MDRO), as well as lack of a written consent, or mental disorders | 621 | 100.0 (621) | 0.0% | PPD ≥4 mm (non-confident) | Full -mouth (four sites—mesial, mid-buccal, mid-lingual, mesial) | 248/373 | 20–80 | NR | NR |
| Zhao et al. (2019) (China) | ≥30 years old | NR | 4930 | 80.2 (3952) | 0.0% | CPI (non-confident) | Partial-mouth (CPI) | 403/3, 549 | 30–68 | NR | NR |
| J. B. Lee et al. (2019) (Republic of Korea) | ≥30 years old | NR | 31,006 | 31.6 (9798) | 0.0% | CPI (non-confident) | Partial-mouth (sextants) | 2771/7027 | ≥30 | 3717/6081 | Company grant |
| Silva Junior et al. (2017) (Brazil) | Piracicaba residents aged 20–64 years old, mentally capable to answer the study questionnaire and agreeing to participate in the research | A physical or psychological state that prevented the achievement of clinical procedures or understanding of the questionnaire | NR | NR | 0.0% | PPD ≥4 mm (non-confident) | Partial-mouth | 43/100 | 20–64 | NR | Research grant |
| | ≥60 years old | NR | 278 | 84.5 (235) | 42.1% | | | 3/47 | ≥60 | NR | NR |

(Continues)

TABLE 1 (Continued)

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response eligibility % (n) | Edentulous participants (n) | Periodontal criteria (confidence) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|---|---|--|--|----------------------------|-----------------------------|--|---|----------------------------|----------------------|--------------|----------------|
| Lasta et al. (2019) (Brazil) | | | | | | CPI (non-confident) | Partial-mouth (sextants) | | | | |
| Botelho et al. (2019) (Portugal) | ≥18 years old (adults and elderly), living in the municipalities of Almada and Seixal | NR | 1064 | 100.0 (1064) | 0.0% | EFP/AAP 2018 | Full-mouth (six sites) | 637/427 | 18–95 | 447/617 | Self-funded |
| Helmi et al. (2019) (Saudi Arabia) | ≥18 years old | Patients who were not within the specified age range, with no BW radiographs, with radiographs in which the cement-enamel junction (CEJ) and alveolar bone crest were not visible, who did not have at least two approximating teeth or where the interproximal space was too narrow to observe the bone crest | 6265 | 18.1 (1131) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (2 sites—mesial and distal) | 893/ 238 | ≥18 | 508/ 623 | Research grant |
| Shimizu et al. (2019) (Japan) | 60–99 years old | Participants without data on carotid intima-media thickness (CIMT) and those without any remaining teeth | 1925 | 47.1 (907) | 0.0% | “ADVANCED” PPD ≥6.0 mm (non-confident) | Full-mouth (2 sites—mesial-buccal and mid-buccal) | 197/710 | 60–99 | 353/NR | Research grant |
| Dhaifullah et al. (2019) (Saudi Arabia) | 18–40 years old At least 20 natural teeth | Pregnancy, periodontal treatment within the past 4 months, a history of systemic diseases or medications known to affect periodontal health status (i.e., diabetes, topical or | 700 | 54.3 (380) | 0.0% | CPI (non-confident) | Partial-mouth (sextant) | 25/283 | 18–40 | 154/154 | Self-funded |

TABLE 1 (Continued)

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response eligibility % (n) | Edentulous participants (n) | Periodontal criteria (confidence) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|--------------------------------------|---|--|--|----------------------------|-----------------------------|-----------------------------------|---|----------------------------|----------------------|--------------|----------------|
| Sun et al. (2020) (China) | 55–74 years old, who had lived longer than 6 months in the sampling areas | Those who withheld consent or had serious diseases | 9054 | 97.0 (8804) | 0.0% | PPD \geq 4 mm (non-confident) | Full-mouth | 6066/2738 | 55–74 | 4514/4540 | Research grant |
| Bongo et al. (2020) (Norway) | Adults in Finnmark County in Northern Norway | Missing questionnaire, missing clinical data, or both, unknown target age, missing written consent, or not accounted for and thus given missing unknown status | 2520 | 82.5 (2078) | 0.0% | EFP/AAP 2018 (confident) | Full-mouth (six sites) | 1032/1046 | 18–75 | 894/1184 | Research grant |
| Romero-Castro et al. (2020) (México) | 18–75 years old, who reside in the state of Guerrero, México | Pregnant or lactating women, as well as patients with systemic diseases or aggressive periodontitis | NR | NR | 0.0% | Armitage 1999 (confident) | Full-mouth | 98/63 | 18–75 | NR | NR |
| Ha et al. (2020) (Australia) | NR | Required antibiotic prophylaxis before dental check-up, congenital heart murmur, heart valve problems, congenital heart disease, bacterial endocarditis, rheumatic fever, kidney disease, haemophilia, pacemaker or automatic defibrillator, hipbone | 4402 | 86 (3792) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (three sites—mesio-buccal, mid-buccal, disto-buccal) | - | \geq 34 | 1854/2011 | Research grant |

(Continues)

TABLE 1 (Continued)

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response participants/ rate/ eligibility % (n) | Edentulous participants (n) | Periodontal criteria (confidence) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|--|---|--|--|--|-----------------------------|-----------------------------------|---|----------------------------|----------------------|--------------|----------------|
| Schmidt et al. (2020) (Switzerland) | ≥55 years old | Edentulous participants; endocarditis | 1673 | 86.9 (1454) | 0.0% | PPD ≥4 mm (non-confident) | Partial-mouth (FDI Index teeth) | 614/840 | ≥55 | NR | Research grant |
| Nakamura et al. (2020) (Japan) | ≥20 years old | Immunocompromised patients (i.e., patients who received chemotherapy, those with severe immunodeficiency, and those with autoimmune disease, who received immunosuppressant therapies) | 201 | 94.5 (190) | 0.0% | PPD ≥4 mm (non-confident) | Full-mouth | 115/75 | ≥20 | NR | Research grant |
| Singh et al. (2020) (Nepal) | 35–44; 65–74 years old | Psychiatric illness | 310 | 100.0 (310) | 0.0% | CPI (non-confident) | NR | 21/96 | 35–44; 65–74 | NR | Self-funded |
| Sekiguchi et al. (2020) (Japan) | ≥18 years old | The participants without all oral examinations, those whose measured probing pocket depth (PPD) and clinical attachment loss (CAL) were not measured, those without General Oral Health Assessment Index (GOHAI) scores, and those who underwent health check-ups more than two times in the 3-year period | 3742 | 31.6 (1183) | 0.0% | PPD ≥4 mm (non-confident) | Full-mouth (two sites- Mesio-Buccal and mid-buccal) | 594/589 | ≥18 | NR | Research grant |
| Diaz-Reissner et al. (2020) (Paraguay) | 18–59 years old, Paraguayans and foreign nationals with more than 14 years of | NR | NR | NR | 1.2% | CPI (non-confident) | Partial-mouth (Ramfjord teeth) | 19/310 | 18–59 | NR | Self-funded |

TABLE 1 (Continued)

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response eligibility % (n) | Edentulous participants (n) | Periodontal criteria (confidence) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|--------------------------------------|--|---|--|----------------------------|-----------------------------|-----------------------------------|----------------------------------|----------------------------|----------------------|--------------|----------------|
| Sakurai et al. (2021) (Japan) | residence in the country Individuals insured by the national health insurance system (including self-employed workers, farmers, and the elderly) and aged 30 years and older | NR | 14,932 | 28.0 (4184) | 0.0% | CPI (non-confident) | Partial-mouth CPITN | 1907/2199 | ≥30 | NR | Research grant |
| Jiao et al. (2021) (China) | 35–45; 55–65; 65–75 years old | NR | 13,463 | 100.0 (13459) | 1.96% | EFP/AAP 2018 (confident) | Full-mouth (most severe site) | 8391/4804 | 35–45; 55–75 | NR | Research grant |
| Goel et al. (2021) (Nepal) | 20–65 years old, tobacco users who were currently consuming tobacco in the form of smoking or smokeless tobacco, non-tobacco users who had never used tobacco in any form (smoke or smokeless tobacco) | Former smokers, patients who actively consume alcohol, patients suffering from known systemic illness, pregnant and lactating females | 1578 | 27.9 (440) | 0.0% | CDC/AAP 2012 (confident) | NR | 315/125 | 20–65 | 186/254 | Self-funded |
| Germeu et al. (2021) (Turkey) | 35–74 years old | Individuals who required antibiotics after routine periodontal procedures | 488 | 488 | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (six sites) | 302/186 | 35–74 | NR | Self-funded |
| Han & Kim (2021) (Republic of Korea) | 40–80 years old | Aged <40 years, edentate, and those missing values in the health assessment or questionnaires | 9450 | 88.1 (8327) | 0.0% | CPI (non-confident) | Partial-mouth (FDI Index teeth) | 3339/4988 | 40–80 | 3572/4755 | Self-funded |
| Costa et al. (2021) (Brazil) | ≥18 years old At least eight natural teeth | People with mental health disorders, under orthodontic treatment, and pregnant women | 450 | 100.0 (450) | 0.0% | CPI (non-confident) | Partial-mouth (sextants) | 304/146 | ≥18 | NR | Research grant |

(Continues)

TABLE 1 (Continued)

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response eligibility % (n) | Edentulous participants (n) | Periodontal criteria (confidence) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|--|--|--|--|----------------------------|-----------------------------|-----------------------------------|--|----------------------------|----------------------|--------------|----------------|
| Song et al. (2021) (Republic of Korea) | Six or more natural teeth | NR | 330 | 91.5 (302) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (2 sites—mesial and distal) | 270/143 | 47–86 | NR | Research grant |
| Gomes-Filho et al. (2021) (Brazil) | >18 years old, registered in basic health units | Diagnosed with neoplasia, HIV, pregnancy, required antibiotic prophylaxis prior to periodontal examination, used anti-inflammatory in the previous 6 months before the examination, received prior periodontal treatment or used antibiotics 6 months before the examination | 1011 | 100.0 (1011) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (6 sites) | 851/160 | ≥18 | 332/NR | Research grant |
| Oliveira et al. (2021) (Brazil) | NR | Requirement of antibiotic prophylactic, psychiatric or mental problems, less than two teeth, less than 18 years old, presenting clinical attachment loss in two adjacent teeth | 1092 | 53.6 (585) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (six sites) | 519/66 | ≥18 | NR | Research grant |
| Clauss et al. (2021) (Burkina Faso) | 35–44 years old, member of a randomly chosen household within the HDSS Household Survey in Nouna | NR | 427 | 100.0 (427) | 0.0% | CPI (non-confident) | Full-mouth (six sites) | 341/86 | 35–44 | NR | Research grant |
| Su et al. (2021) (Japan) | Smokers and patients with a medical history of hypertension, | Subjects with oral cancer or potentially malignant oral disorders (i.e., | 149 | 83.2 (124) | 0.0% | PPD ≥4 mm (non-confident) | Full-mouth (6 sites) | 72/52 | 35–90 | NR | Research grant |

TABLE 1 (Continued)

| Authors (year) (country) | Inclusion criteria | Exclusion criteria | Invited participants/ initial sample (n) | Response participants/ rate/ eligibility % (n) | Edentulous participants (n) | Periodontal criteria (confidence) | Periodontal examination protocol | Periodontitis/ healthy (n) | Age interval (years) | Male/ female | Funding source |
|------------------------------------|--|---|--|--|-----------------------------|-----------------------------------|----------------------------------|----------------------------|----------------------|--------------|----------------|
| | diabetes, hyperlipidemia, stroke, heart disease, or bone and joint disease | leukoplakia or lichen planus), cancer patients receiving surgical treatment, chemotherapy or radiotherapy, those with auto-immune diseases receiving steroid therapy and those with severe immunodeficiency | | | | | | | | | |
| Iwasaki et al. (2021) (Japan) | At least 18 years of age and able to read and understand Japanese | Having fewer than two teeth and previous diagnosis of a severe or terminal disease, such as advanced heart failure, end-stage kidney disease, or advanced-stage cancer | 215 | 92.6 (199) | 0.0% | CDC/AAP 2012 (confident) | Full-mouth (six sites) | 118/81 | 19–77 | NR | Research grant |
| Stødle et al. (2021) (Norway) | ≥19 years old and residents of the county | No radiographs, edentulous | 7347 | 67.1 (4933) | 0.66% | EFP/AAP 2018 (confident) | Full-mouth (six sites) | 3.573/1.290 | ≥19 | NR | Research grant |
| Bigün Çetin et al. (2021) (Turkey) | NR | Patients younger than 18 years, edentulous patients | 573 | 94.4 (541) | 0.0% | EFP/AAP 2018 (confident) | Full-mouth (six sites) | 221/320 | ≥18 | 252/289 | Research grant |
| Kocher et al. (2021) (Germany) | Adults 35–44; 65–74 | NR | 2007 | 92.2 (1851) | 0.0% | CPI (non-confident) | Partial-mouth (FDI Index teeth) | 893/948 | 35–44; 65–74 | NR | Research grant |

Abbreviations: AAP, American Academy of Periodontology; CAL, clinical attachment level; CDC, Centers for Disease Control; CPI, Community Periodontal Index; EFP, European Federation of Periodontology; NR, not reported; PPD, periodontal probing depth.

TABLE 2 Periodontitis prevalence according to the type of confident case definition

| | <i>n</i> | ES | 95% CI | <i>T</i> | <i>T</i> ² | <i>I</i> ² (%) | <i>p</i> -value | Egger test [SE] (<i>p</i> -value) |
|------------------------------|----------|------|-----------|----------|-----------------------|---------------------------|-----------------|------------------------------------|
| Overall | 24 | 61.6 | 55.1–67.9 | 0.16 | 0.03 | 99.1 | <.000001 | 0.61 [3.63] (.867) |
| Risk of bias | | | | | | | | |
| Low | 7 | 54.1 | 47.4–60.6 | 0.09 | 0.01 | 98.9 | <.000001 | - |
| Moderate | 17 | 64.7 | 56.3–72.6 | 0.18 | 0.03 | 99.0 | <.000001 | -3.76 [5.58] (.511) |
| Case definition | | | | | | | | |
| EFP/AAP 2018 | 6 | 52.3 | 38.2–66.2 | 0.18 | 0.03 | 99.4 | <.000001 | - |
| CDC/AAP 2012 | 12 | 68.1 | 59.7–76.0 | 0.15 | 0.02 | 99.2 | <.000001 | 12.35 [5.20] (.039) |
| CDC/AAP 2007 | 3 | 48.8 | 38.2–66.2 | 0.10 | 0.01 | 95.0 | <.000001 | - |
| AAP 1999 | 3 | 66.7 | 61.3–71.8 | 0.04 | 0.00 | 50.5 | <.000001 | - |
| Age interval | | | | | | | | |
| Wider age interval (≥18/≥19) | 11 | 57.1 | 45.9–67.9 | 0.19 | 0.04 | 99.4 | <.000001 | -9.31 [7.73] (.259) |
| ≥30 years old | 7 | 61.4 | 55.4–67.2 | 0.08 | 0.01 | 99.3 | <.000001 | - |
| <65 years old | 2 | 59.4 | 34.5–81.9 | 0.18 | 0.03 | 98.7 | <.000001 | - |
| ≥65 years old | 3 | 79.3 | 64.2–91.2 | 0.15 | 0.02 | 97.5 | <.000001 | - |
| Continent | | | | | | | | |
| Asia | 10 | 62.4 | 55.0–69.5 | 0.15 | 0.01 | 97.6 | <.000001 | -0.91 [3.16] (.780) |
| Europe | 5 | 65.5 | 48.7–80.5 | 0.19 | 0.04 | 99.5 | <.000001 | - |
| North America | 4 | 62.6 | 48.3–75.9 | 0.15 | 0.02 | 99.2 | <.000001 | - |
| South America | 5 | 54.9 | 33.4–75.5 | 0.06 | 0.25 | 99.4 | <.000001 | - |

Note: Effect size (ES) with 95% confidence interval (CI), tau (*T*), tau squared (*T*²) and *I*-squared (*I*²), *p*-value and Egger test are presented.

intervals defined in the studies ($p = .0101$). Yet, studies reporting prevalence data of elderly participants (≥65 years or older) presented the highest estimates with an average a 79.3% overall percentage (95% CI: 64.2–91.2, $p < .000001$, $I^2 = 97.5\%$).

3.4.2 | Prevalence per continent

Considering the geographic location, the pooled estimate from the European continent was the highest (65.5%), while the South American continent had the lowest (54.9%; Table 2). Yet, all continents reported significant pooled estimates exceeding 50%.

3.4.3 | Severity

Concerning severity (Table 3), the pooled measures pointed to an estimated moderate to severe periodontitis prevalence of 53.2% (95% CI: 44.3–61.9, $p < .000001$, $I^2 = 99.2\%$). Although having substantial heterogeneity, this estimate was relatively stable and not influenced by the female/male ratio (-0.220 , $p = .2952$), latitude (0.001 , $p = .3641$), longitude (0.000 , $p = .8570$), or sample size (-0.000 , $p = .8172$). The estimated prevalence of severe periodontitis was 23.6% (95% CI: 17.6–30.1, $p < .000001$, $I^2 = 99.4\%$). Similarly, we observed high heterogeneity, and this estimate was significantly influenced by the male/female ratio (-0.401 , $p = .0334$) but not by the remaining variables (sample size, latitude, and longitude).

3.5 | Additional analysis

When inspecting the existence of publication bias, funnel asymmetry was found in studies reporting periodontitis using the CDC/AAP 2012 (estimate = 12.35, SE = 5.204, $p = .039$; Table 2). For the remaining estimates, no publication bias was observed.

4 | DISCUSSION

4.1 | Summary of main results

The results of this systematic review confirm that periodontitis continues to be an alarming world public health problem, with nearly 62% pooled prevalence in dentate adults as seen from studies performed between 2011 and 2020. Pooled prevalence for moderate to severe cases was 53.2% while for severe periodontitis it was 23.6%.

Pooled estimates from confident case definitions presented almost double the prevalence. The CDC/AAP 2012 classification was the most used diagnostic criterion and gave the highest prevalence estimate (68.1%), while the EFP/AAP 2018 case definition gave a pooled estimate of 52.3%.

4.2 | Evidence quality and potential bias in the review process

This study's strengths and limitations are based primarily on the methodological quality of the included studies per se, while the study was

TABLE 3 Periodontitis prevalence according to the form of periodontitis

| Periodontitis | n | ES | 95% CI | T | T ² | I ² (%) | p-value | Egger test [SE] (p-value) |
|--------------------------------------|----|------|-----------|------|----------------|--------------------|----------|---------------------------|
| Moderate to severe versus mild to no | 20 | 53.2 | 44.3–61.9 | 0.20 | 0.04 | 99.2 | <.000001 | 2.38 [4.32] (.589) |
| Severe versus non-severe | 20 | 23.6 | 17.6–30.1 | 0.17 | 0.03 | 99.4 | <.000001 | 1.29 [5.00] (.799) |

Note: Effect size (ES) with 95% confidence interval (CI), tau (T), tau squared (T²) and I-squared (I²), p-value, and Egger test are presented.

prepared and reported according to state-of-the-art guidelines and comprehensively analysed the possible confounding variables that could affect the results. Indeed, prevalence was strictly affected by periodontal case definitions with regard to their confidence. As for the remaining variables, they did not influence the final estimates. Defining the confidence of a case definition always implies an uncertain degree of selection bias, yet this decision was based on a previous study by Muñoz-Aguilera (2020), which found a significant weight on the results. Similarly, studies with confident case definitions reported almost twice the prevalence as those using non-confident classifications according to the present results.

Still on the periodontal case definitions, our meta-analytical estimates differ from those of previous studies using the GBD approach (Marcenes et al., 2013; Kassebaum et al., 2014; Chen et al., 2021; Wu et al., 2022) and aligns with a recent meta-analysis on the global prevalence of aggressive periodontitis (Bouziane et al., 2020). On one hand, GBD estimates are based on complex and strong iterative approaches and thus provide worldwide approximations (IHME, 2018) yet using less reliable diagnostic criteria (such as the CPI of treatment needs). On the other hand, our approach provides more reliable estimations based on full-mouth examinations and diagnostic consensus; nevertheless, the existence of several confident case definitions confuses the comparison among studies. Consequently, arriving at a consensus to strengthen the surveillance of periodontitis will certainly benefit periodontal research and future meta-epidemiological studies. Defining mandatory periodontal clinical measures (such as PPD, CAL, PISA, or PESA) and thresholds and providing results based on at least two diagnostic criteria (CDC/AAP 2012 and EFP/AAP 2018) are suggestions that may contribute to the robustness of periodontal prevalence reporting. Yet, narrowing the results according to the confidence of case definition may have certainly resulted in less geographic coverage and was contingent upon available epidemiological studies. Only 25 countries (17 of which were using confident diagnostic criteria) were accounted in the final analyses, without representation from Africa and Oceania.

One possible explanation for the obtained differences in results between the 2007 definition and its updated version of 2012 may be due to the fact the original version was designed to capture only moderate and severe periodontitis while in the 2012 update the mild case was introduced. The incorporation of a mild stage may have reduced the number of periodontally healthy cases, thereby leading to a potential increase in the number of periodontitis cases, a factor that could partially explain these noticeable differences.

Finally, several studies were lacking in information on male/female prevalence ratio, clear age intervals, distribution according to

smoking habits, socio-economic data, and distribution of periodontitis according to its severity. Thus, our examination on additional sources of heterogeneity and interpretation of the results was restricted and could be expanded in the future if studies provide such information. Moreover, the inclusion of edentulous participants may have an impact on the prevalence reported, but due to missing data, we were not able to measure its magnitude.

4.3 | Agreements and disagreements with other reviews or studies

There is overall agreement that the prevalence and incidence of periodontitis have been increasing, possibly due to population growth and ageing (Marcenes et al., 2013; Kassebaum et al., 2017; Peres et al., 2019; Watt et al., 2019; GBD 2017 Oral Disorders Collaborators et al., 2020; Wu et al., 2022). These results align with this consensus and highlight the need for better periodontal care programmes mainly focusing the poor and socially disadvantaged populations that are disproportionately affected. As such, these estimates can flag worrisome levels that will contribute to the World Health Organization Resolution on Oral Health within the road map for neglected tropical diseases for 2021–2030 (WHO, 2021).

In Wu et al. (2022), GBD data allowed the assessment of all regions and countries, which was not possible with our estimates. In the latter, periodontitis was mostly prevalent in 55–59-year-olds, with substantial prevalence from 55 years onwards. Similarly, our age subgroup meta-analysis showed individuals of 65 years or older to have the highest prevalence when compared with younger age groups. Regarding severe periodontitis, our results report a substantial prevalence of severe periodontitis (24%). Compared to the 11.2% of severe periodontitis reported globally by Kassebaum et al. (2014), our estimate is definitely higher. However, the methodologies used in the two estimates differ, as in Kassebaum et al. the results relied on probing depth, which leads to an underestimation of periodontitis in adults. Despite this estimation discrepancy, our estimates point to a worrisome forecast of severe periodontitis among dentate adults.

The findings from this study can inform global health stakeholders about the prevalence of periodontitis from epidemiological studies to preempt and better manage public periodontal health strategies. However, because of the limited number of countries represented in the final sample, the generalizability of these estimates is limited. Therefore, we emphasize the need for continued epidemiological surveillance, from both national and regional settings, using appropriate

diagnostic strategies, to better convey accurate estimates and to allow meta-analytical global outputs.

5 | CONCLUSION

Within the limits of this systematic review, it can be concluded that, over the last decade (from 2011 to 2020), the estimated pooled prevalence of periodontitis is nearly 60%, with its severe stage affecting approximately 24% of the studied population. These results show an alarmingly high prevalence compared to estimates from 1990 to 2010.

Based on this data, the worldwide periodontitis estimates of this study are crucial to establish priorities for research, development, public health policy, and funding to governments and domestic and international non-governmental organizations.

AUTHOR CONTRIBUTIONS

Diogo Trindade contributed to data acquisition and interpretation and drafted and critically revised the manuscript; João Botelho contributed to conception, design, data acquisition, and analysis and drafted and critically revised the manuscript; Vanessa Machado contributed to conception and design and critically revised the manuscript; Rui Carvalho contributed to data interpretation and critically revised the manuscript; José João Mendes and Leandro Chambrone contributed to data analysis and interpretation and critically revised the manuscript. All authors gave final approval and agreed to be accountable for all aspects of the work.

FUNDING INFORMATION

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

The data that supports the findings of this study are available in the supplementary material of this article.

ORCID

Rui Carvalho  <https://orcid.org/0000-0002-1429-3073>

Vanessa Machado  <https://orcid.org/0000-0003-2503-260X>

Leandro Chambrone  <https://orcid.org/0000-0002-2838-1015>

José João Mendes  <https://orcid.org/0000-0003-0167-4077>

João Botelho  <https://orcid.org/0000-0002-1019-8263>

REFERENCES

Araya Vallespir, C., Ulloa Ortega, C., Luengo Machuca, L., Rodríguez Vera, M., & Contreras, S. S. (2014). Grado de concordancia de los índices más utilizados en estudios epidemiológicos de la enfermedad periodontal. *Revista Clínica de Periodoncia, Implantología y Rehabilitación Oral*, 7(3), 175–179.

- Armitage, G. C. (1999). Development of a classification system for periodontal diseases and conditions. *Annals of Periodontology*, 4(1), 1–6.
- Australian Research Centre for Population Oral Health. (2019). *Australia's Oral Health National Study of Adult Oral Health 2017–18*. The University of Adelaide.
- Balaji, S., Lavu, V., & Rao, S. (2018). Chronic periodontitis prevalence and the inflammatory burden in a sample population from South India. *Indian Journal of Dental Research*, 29(2), 254.
- Barendregt, J. J., Doi, S. A., Lee, Y. Y., Norman, R. E., & Vos, T. (2013). Meta-analysis of prevalence. *Journal of Epidemiology and Community Health*, 67(11), 974–978.
- Belbasis, L., & Bellou, V. (2018). Introduction to epidemiological studies. In E. Evangelou (Ed.), *Genetic epidemiology (Methods in Molecular Biology)* (Vol. 1793, pp. 1–6). Springer New York. https://doi.org/10.1007/978-1-4939-7868-7_1
- Bhat, M., Do, L., & Roberts-Thomson, K. (2018). Risk indicators for prevalence, extent and severity of periodontitis among rural Indian population aged 35–54 years. *International Journal of Dental Hygiene*, 16(4), 492–502.
- Bhat, M., Roberts-Thomson, K., & Do, L. G. (2015). Clustering of risk indicators for periodontal disease: A population-based study. *Community Dental Health*, 32(3), 158–162.
- Bilgin Çetin, M., Sezgin, Y., Maraş, E., & Cebeci, İ. A. (2021). Association of probable bruxism with periodontal status: A cross-sectional study in patients seeking periodontal care. *Journal of Periodontal Research*, 56(2), 370–378.
- Bongo, A. K. S., Brustad, M., Oscarson, N., & Jönsson, B. (2020). Periodontal health in an indigenous Sámi population in northern Norway: A cross-sectional study. *BMC Oral Health*, 20(1), 104.
- Botelho, J., Machado, V., Leira, Y., Proença, L., Chambrone, L., & Mendes, J. J. (2022). Economic burden of periodontitis in the United States and Europe—An updated estimation. *Journal of Periodontology*, 93(3), 373–379.
- Botelho, J., Machado, V., Proença, L., Alves, R., Cavacas, M. A., Amaro, L., & Mendes, J. J. (2019). Study of periodontal health in Almada-Seixal (SoPHIAS): A cross-sectional study in the Lisbon metropolitan area. *Scientific Reports*, 9(1), 15538.
- Botelho, J., Machado, V., Proença, L., Bellini, D. H., Chambrone, L., Alcoforado, G., & Mendes, J. J. (2020). The impact of nonsurgical periodontal treatment on oral health-related quality of life: A systematic review and meta-analysis. *Clinical Oral Investigations*, 24(2), 585–596.
- Botelho, J., Mascarenhas, P., Viana, J. et al. (2022). An umbrella review of the evidence linking oral health and systemic noncommunicable diseases. *Nature Communications*, 13, 7614. <https://doi.org/10.1038/s41467-022-35337-8>
- Bouziane, A., Hamdoun, R., Abouqal, R., & Ennibi, O. (2020). Global prevalence of aggressive periodontitis: A systematic review and meta-analysis. *Journal of Clinical Periodontology*, 47(4), 406–428.
- Buset, S. L., Walter, C., Friedmann, A., Weiger, R., Borgnakke, W. S., & Zitzmann, N. U. (2016). Are periodontal diseases really silent? A systematic review of their effect on quality of life. *Journal of Clinical Periodontology*, 43(4), 333–344.
- Caton, J. G., Armitage, G., Berglundh, T., Chapple, I. L. C., Jepsen, S., Kornman, K. S., Mealey, B. L., Papapanou, P. N., Sanz, M., & Tonetti, M. S. (2018). A new classification scheme for periodontal and peri-implant diseases and conditions—Introduction and key changes from the 1999 classification. *Journal of Periodontology*, 89(Suppl 1), S1–S8.
- Chen, M. X., Zhong, Y. J., Dong, Q. Q., Wong, H. M., & Wen, Y. F. (2021). Global, regional, and national burden of severe periodontitis, 1990–2019: An analysis of the Global Burden of Disease Study 2019. *Journal of Clinical Periodontology*, 48(9), 1165–1188.
- Clauss, A., Sie, A., Zabre, P., Schmoll, J., Sauerborn, R., & Listl, S. (2021). Population-based prevalence of oral conditions as a basis for planning

- community-based interventions: An epidemiological study from rural Burkina Faso. *Frontiers in Public Health*, 9, 697498.
- Costa, P., Canaan, J., Midori Castelo, P., Campideli Fonseca, D., Márcia Pereira-Dourado, S., Mendonça Murata, R., Pardi, V., & Pereira, L. J. (2021). Influence of micronutrient intake, sociodemographic, and behavioral factors on periodontal status of adults assisted by a public health care system in Brazil: A cross-sectional multivariate analysis. *Nutrients*, 13(3), 973.
- Dhaifullah, E., Al-Maweri, S. A., Koppolu, P., Elkhat, E., Mostafa, D., & Mahgoub, M. (2019). Body mass index and periodontal health status among young Saudi adults: A cross-sectional study. *Annals of Saudi Medicine*, 39(6), 433–440.
- Díaz-Reissner, C., Roldán-Merino, J., & Casas-García, I. (2020). Association between self-assessment and clinical oral health status in adults, Paraguay. *Journal of Oral Research*, 9(6), 483–489.
- Eke, P. I., Dye, B. A., Wei, L., Slade, G. D., Thornton-Evans, G. O., Borgnakke, W. S., Taylor, G. W., Page, R. C., Beck, J. D., & Genco, R. J. (2015). Update on prevalence of periodontitis in adults in the United States: NHANES 2009 to 2012. *Journal of Periodontology*, 86(5), 611–622.
- Eke, P. I., Page, R. C., Wei, L., Thornton-Evans, G., & Genco, R. J. (2012). Update of the case definitions for population-based surveillance of periodontitis. *Journal of Periodontology*, 83(12), 1449–1454.
- Eke, P. I., Thornton-Evans, G. O., Wei, L., Borgnakke, W. S., Dye, B. A., & Genco, R. J. (2018). Periodontitis in US adults. *The Journal of the American Dental Association*, 149(7), 576–588.e6.
- Figueiredo, A., Soares, S., Lopes, H., dos Santos, J. N., Ramalho, L. M. P., Cangussu, M. C., & Cury, P. R. (2013). Destructive periodontal disease in adult Indians from Northeast Brazil: Cross-sectional study of prevalence and risk indicators. *Journal of Clinical Periodontology*, 40(11), 1001–1006.
- Frencken, J. E., Sharma, P., Stenhouse, L., Green, D., Laverty, D., & Dietrich, T. (2017). Global epidemiology of dental caries and severe periodontitis—A comprehensive review. *Journal of Clinical Periodontology*, 44(Suppl 18), S94–S105.
- GBD 2017 Oral Disorders Collaborators, Bernabe, E., Marcenes, W., Hernandez, C. R., Bailey, J., Abreu, L. G., Alipour, V., Amini, S., Arabloo, J., Arefi, Z., & Arora, A. (2020). Global, regional, and national levels and trends in burden of oral conditions from 1990 to 2017: A systematic analysis for the global burden of disease 2017 study. *Journal of Dental Research*, 99(4), 362–373.
- Germen, M., Baser, U., Lacin, C. C., Firatli, E., İşsever, H., & Yalcin, F. (2021). Periodontitis prevalence, severity, and risk factors: A comparison of the AAP/CDC case definition and the EFP/AAP classification. *IJERPH*, 18(7), 3459.
- Ghassib, I. H., Batarseh, F. A., Wang, H., & Borgnakke, W. S. (2021 Aug). Clustering by periodontitis-associated factors: A novel application to NHANES data. *Journal of Periodontology*, 92(8), 1136–1150.
- Giacaman, R. A., Sandoval Salas, D., Bustos Alvarez, I. P., Rojas Cáceres, M. A., & Mariño, R. J. (2016). Epidemiología del estado de salud periodontal en la VII Región del Maule, Chile. *Revista Clínica de Periodoncia, Implantología y Rehabilitación Oral*, 9(2), 184–192.
- Goel, K., Sharma, S., Baral, D. D., & Agrawal, S. K. (2021). Current status of periodontitis and its association with tobacco use amongst adult population of Sunsari district, in Nepal. *BMC Oral Health*, 21(1), 66.
- Gomes-Filho, I. S., Santos, P. N. P., Cruz, S. S., Figueiredo, A. C. M. G., Trindade, S. C., Ladeia, A. M., Cerqueira, E. M., Passos-Soares, J. S., Coelho, J. M., Hintz, A. M., & Barreto, M. L. (2021). Periodontitis and its higher levels of severity are associated with the triglyceride/high density lipoprotein cholesterol ratio. *Journal of Periodontology*, 92(11), 1509–1521.
- Ha, D. H., John Spencer, A., Ju, X., & Do, L. G. (2020). Periodontal diseases in the Australian adult population. *Australian Dental Journal*, 65, S52–S58. <https://doi.org/10.1111/adj.12765>
- Han, D. H., & Kim, M. S. (2021). Are occupational and environmental noises associated with periodontitis? Evidence from a Korean representative cross-sectional study. *BMC Public Health*, 21(1), 616.
- He, S., Wei, S., Wang, J., & Ji, P. (2018). Chronic periodontitis and oral health-related quality of life in Chinese adults: A population-based, cross-sectional study. *Journal of Periodontology*, 89(3), 275–284.
- Helmi, M. F., Huang, H., Goodson, J. M., Hasturk, H., Tavares, M., & Natto, Z. S. (2019). Prevalence of periodontitis and alveolar bone loss in a patient population at Harvard School of Dental Medicine. *BMC Oral Health*, 19(1), 254.
- Higgins, J. P., Thomas, J., Chandler, J., Cumpston, M., Li, T., Page, M. J., & Welch, V. A. (2019). *Cochrane Handbook for Systematic Reviews of Interventions* (2nd ed.). Wiley <https://www.wiley.com/en-gb/Cochrane+Handbook+for+Systematic+Reviews+of+Interventions%2C+2nd+Edition-p-9781119536628>.
- Holde, G. E., Oscarson, N., Trovik, T. A., Tillberg, A., & Jönsson, B. (2017). Periodontitis prevalence and severity in adults: A cross-sectional study in Norwegian circumpolar communities. *Journal of Periodontology*, 88(10), 1012–1022.
- Holtfreter, B., Albandar, J. M., Dietrich, T., Dye, B. A., Eaton, K. A., Eke, P. I., Papapanou, P. N., & Kocher, T. (2015). Standards for reporting chronic periodontitis prevalence and severity in epidemiologic studies: Proposed standards from the Joint EU/USA Periodontal Epidemiology Working Group. *Journal of Clinical Periodontology*, 42(5), 407–412.
- Hoy, D., Brooks, P., Woolf, A., Blyth, F., March, L., Bain, C., Baker, P., Smith, E., & Buchbinder, R. (2012). Assessing risk of bias in prevalence studies: Modification of an existing tool and evidence of interrater agreement. *Journal of Clinical Epidemiology*, 65(9), 934–939.
- IHME. (2018). *Protocol for the global burden of diseases, injuries, and risk factors study (GBD)*. Institute for Health Metrics and Evaluation.
- Iwasaki, M., Usui, M., Ariyoshi, W., Nakashima, K., Nagai-Yoshioka, Y., Inoue, M., Kobayashi, K., & Nishihara, T. (2021). Interruption of regular dental visits during the COVID-19 pandemic due to concerns regarding dental visits was associated with periodontitis in Japanese office workers. *Journal of Periodontal Research*, 56(6), 1091–1098.
- Iwasaki, T., Hirose, A., Azuma, T., Ohashi, T., Watanabe, K., Obora, A., Deguchi, F., Kojima, T., Isozaki, A., & Tomofuji, T. (2018). Correlation between ultrasound-diagnosed non-alcoholic fatty liver and periodontal condition in a cross-sectional study in Japan. *Scientific Reports*, 8(1), 7496.
- Jaafar, N., Hakim, H., Mohd Nor, N. A., Mohamed, A., Saub, R., Esa, R., Doss, J., Mohd Yusof, Z. Y., Ab-Murat, N., Abu Kassim, N. L., & Majid, H. A. (2014). Is the burden of oral diseases higher in urban disadvantaged community compared to the national prevalence? *BMC Public Health*, 14(S3), S2.
- Jiao, J., Jing, W., Si, Y., Feng, X., Tai, B., Hu, D., Lin, H., Wang, B., Wang, C., Zheng, S., & Liu, X. (2021). The prevalence and severity of periodontal disease in mainland China: Data from the fourth National Oral Health Survey (2015–2016). *Journal of Clinical Periodontology*, 48(2), 168–179.
- Juarez, I., Hernández, M., Letelier, C., Halabí, D., & Araneda, C. (2014). Association between abdominal obesity and periodontal disease. Cross-sectional study. *Journal of Oral Research*, 4(2), 95–102.
- Kassebaum, N. J., Bernabé, E., Dahiya, M., Bhandari, B., Murray, C. J. L., & Marcenes, W. (2014). Global burden of severe periodontitis in 1990–2010: A systematic review and meta-regression. *Journal of Dental Research*, 93(11), 1045–1053.
- Kassebaum, N. J., Smith, A. G. C., Bernabé, E., Fleming, T. D., Reynolds, A. E., Vos, T., Murray, C. J., Marcenes, W., & GBD 2015 Oral Health Collaborators. (2017). Global, regional, and national prevalence, incidence, and disability-adjusted life years for oral conditions for 195 countries, 1990–2015: A systematic analysis for the global burden of diseases, injuries, and risk factors. *Journal of Dental Research*, 96(4), 380–387.

- Khan, S., Khalid, T., & Awan, K. H. (2016). Chronic periodontitis and smoking. Prevalence and dose-response relationship. *Saudi Medical Journal*, 37(8), 889–894.
- Kocher, T., Holtfreter, B., Pitchika, V., Kuhr, K., & Jordan, R. (2021). Entwicklung der Zahn- und Mundgesundheit in Deutschland von 1997 bis 2014. *Bundesgesundheitsbl.*, 64, 782–792.
- Lasta, R., Devilla, A., Simoni, V., Foralosso, J. C., Kellermann, M. G., Marcon, S., Corralo, V. D., & Rodrigues-Junior, S. A. (2019). Oral health profile of participants of an elderly cohabitation program in the state of Santa Catarina, Brazil. *Revista de Odontologia da UNESP*, 48, e20190061.
- Lee, J. B., Shin, H. J., Kim, D. Y., & Pang, E. K. (2019). Evaluation of prognosis related to compliance with supportive periodontal treatment in patients with chronic periodontitis: A clinical retrospective study. *Journal of Periodontal & Implant Science*, 49(2), 76.
- Lee, K., & Kim, J. (2019). Dairy food consumption is inversely associated with the prevalence of periodontal disease in Korean adults. *Nutrients*, 11(5), 1035.
- Marcenes, W., Kassebaum, N. J., Bernabé, E., Flaxman, A., Naghavi, M., Lopez, A., & Murray, C. J. (2013). Global burden of oral conditions in 1990–2010: A systematic analysis. *Journal of Dental Research*, 92(7), 592–597.
- Moya, P., Chappuzeau, E., Caro, J. C., & Monsalves, M. J. (2012). Situación de salud oral y calidad de vida de los adultos mayores. *Revista Estomatológica Herediana*, 22(4), 197–202.
- Muñoz Aguilera, E., Suvan, J., Buti, J., Czesnikiewicz-Guzik, M., Barbosa Ribeiro, A., Orlandi, M., Guzik, T. J., Hingorani, A. D., Nart, J., & D'Aiuto, F. (2020). Periodontitis is associated with hypertension: A systematic review and meta-analysis. *Cardiovascular Research*, 116(1), 28–39.
- Nakamura, M., Shigeishi, H., Cheng-Yih, S., Sugiyama, M., & Ohta, K. (2020). Oral human cytomegalovirus prevalence and its relationships with periodontitis and *Porphyromonas gingivalis* in Japanese adults: A cross-sectional study. *Journal of Applied Oral Science*, 28, e20200501.
- Oliveira, L. M., Cerezer, D. M., Casarin, M., Moreira, C. H. C., & Zanatta, F. B. (2021). Alcohol use disorders are associated with higher prevalence of periodontitis in a rural area of Brazil. *Journal of Periodontal Research*, 56(5), 940–950.
- Ortiz, A. P., González, D., Ramos, J., Muñoz, C., Reyes, J. C., & Pérez, C. M. (2018). Association of marijuana use with oral HPV infection and periodontitis among Hispanic adults: Implications for oral cancer prevention. *Journal of Periodontology*, 89(5), 540–548.
- Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C. D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., & Chou, R. (2021). The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ*, 29, n71. <https://doi.org/10.1136/bmj.n71>
- Page, R. C., & Eke, P. I. (2007). Case definitions for use in population-based surveillance of periodontitis. *Journal of Periodontology*, 78(7 s), 1387–1399.
- Peres, M. A., Macpherson, L. M. D., Weyant, R. J., Daly, B., Venturelli, R., Mathur, M. R., Listl, S., Celeste, R. K., Guarnizo-Herreño, C. C., Kearns, C., & Benzian, H. (2019). Oral diseases: A global public health challenge. *The Lancet*, 394(10194), 249–260.
- Petersen, P. E. (2009). Global policy for improvement of oral health in the 21st century—Implications to oral health research of World Health Assembly 2007, World Health Organization. *Community Dentistry and Oral Epidemiology*, 37(1), 1–8.
- Pinto-Filho, J. M., Ribeiro, L. S. F., Sartori, L., dos Santos, J. N., Ramalho, L. M. P., & Cury, P. R. (2018). Association between alcohol dependence and both periodontal disease and tooth loss: A cross-sectional study. *Environmental Science and Pollution Research*, 25(29), 29089–29095.
- Ramírez, J. C. M., Lopera, N. S., López, A. P., Agudelo-Suárez, A. A., & Botero, J. E. (2017). Periodontal disease and its relationship with clinical and sociodemographic variables in adult patients treated in a service/teaching institution. *Revista Odontológica Mexicana*, 21(3), 160–167.
- Romero-Castro, N. S., Castro-Alarcón, N., Reyes-Fernández, S., Flores-Alfaro, E., Serna-Radilla, V. O., & Parra-Rojas, I. (2020). Periodontal disease distribution, risk factors, and importance of primary healthcare in the clinical parameters improvement. *International Journal of Odontostomatology*, 14(2), 183–190.
- Sakurai, A., Yamada, S., Karasawa, I., Kondo, E., & Kurita, H. (2021). Accuracy of a salivary examination kit for the screening of periodontal disease in a group medical check-up (Japanese-specific health check-up). *Medicine*, 100(6), e24539.
- Schmidt, J. C., Vogt, S., Imboden, M., Schaffner, E., Grize, L., Zemp, E., Probst-Hensch, N., & Zitzmann, N. U. (2020). Dental and periodontal health in a Swiss population-based sample of older adults: A cross-sectional study. *European Journal of Oral Sciences*, 128(6), 508–517.
- Schünemann, H. J., Oxman, A. D., Brozek, J., Glasziou, P., Jaeschke, R., Vist, G. E., Williams, J. W., Kunz, R., Craig, J., Montori, V. M., & Bossuyt, P. (2008). Grading quality of evidence and strength of recommendations for diagnostic tests and strategies. *BMJ*, 336(7653), 1106–1110.
- Schwarzer, G. (2007). Meta: An R package for meta-analysis. *R News*, 7, 40–45.
- Schwarzer, G., Carpenter, J. R., & Rücker, G. (2015). *Meta-analysis with R*. Springer.
- Schwendicke, F., Dörfer, C. E., & Meier, T. (2018). Global smoking-attributable burden of periodontal disease in 186 countries in the year 2015. *Journal of Clinical Periodontology*, 45(1), 2–14.
- Sekiguchi, A., Kawashiri, S., Hayashida, H., Nagaura, Y., Nobusue, K., Nonaka, F., Yamanashi, H., Kitamura, M., Kawasaki, K., Fukuda, H., & Iwasaki, T. (2020). Association between high psychological distress and poor oral health-related quality of life (OHQoL) in Japanese community-dwelling people: The Nagasaki Islands Study. *Environmental Health and Preventive Medicine*, 25(1), 82.
- Shariff, J. A., Burkett, S., Watson, C. W. -M., Cheng, B., Noble, J. M., & Papananou, P. N. (2018). Periodontal status among elderly inhabitants of northern Manhattan: The WHICAP ancillary study of oral health. *Journal of Clinical Periodontology*, 45(8), 909–919.
- Shimizu, Y., Yamanashi, H., Kitamura, M., Furugen, R., Iwasaki, T., Fukuda, H., Hayashida, H., Kawasaki, K., Kiyoura, K., Kawashiri, S. Y., & Saito, T. (2019). Association between human T cell leukemia virus type-1 (HTLV-1) infection and advanced periodontitis in relation to atherosclerosis among elderly Japanese: A cross-sectional study. *Environmental Health and Preventive Medicine*, 24(1), 81.
- Shyagali, T., Salama, M. H., & Bhayya, D. (2017). Prevalence of tobacco usage and its effect on the periodontal health parameters in the mining employees and the general population—A comparative study. *Brazilian Journal of Oral Science*, 16, 1–9.
- Silva-Junior, M. F., de Sousa, A. C. C., Batista, M. J., & da LR de Sousa, M. (2017). Condição de saúde bucal e motivos para extração dentária entre uma população de adultos (20-64 anos). *Ciênc Saúde Coletiva*, 22(8), 2693–2702.
- Singh, A., Shrestha, A., Bhagat, T. K., & Baral, D. D. (2020). Assessment of oral health status and treatment needs among people of Foklyan area, Dharan, Nepal. *BMC Oral Health*, 20(1), 320.
- Skośkiewicz-Malinowska, K., Malicka, B., Ziętek, M., & Kaczmarek, U. (2018). Oral health condition and occurrence of depression in the elderly. *Medicine*, 97(41), e12490.
- Song, E., Park, M. J., Kim, J. A., Roh, E., Yu, J. H., Kim, N. H., Yoo, H. J., Seo, J. A., Kim, S. G., Kim, N. H., & Baik, S. H. (2021). Implication of thyroid function in periodontitis: A nationwide population-based study. *Scientific Reports*, 11(1), 22127.
- Stødle, I. H., Verket, A., Høvik, H., Sen, A., & Koldsland, O. C. (2021). Prevalence of periodontitis based on the 2017 classification in a Norwegian population: The HUNT study. *Journal of Clinical Periodontology*, 48(9), 1189–1199.

- Su, C. Y., Shigeishi, H., Murodumi, H., Sugiyama, M., Ohta, K., & Takemoto, T. (2021). Association of oral Epstein-Barr virus with periodontal health in Japanese adults. *Experimental and Therapeutic Medicine*, 22(1), 767.
- Sun, H., Du, M., Tai, B., Chang, S., Wang, Y., & Jiang, H. (2020). Prevalence and associated factors of periodontal conditions among 55- to 74-year-old adults in China: Results from the 4th National Oral Health Survey. *Clinical Oral Investigations*, 24(12), 4403–4412.
- Thanakun, S., Watanabe, H., Thaweboon, S., & Izumi, Y. (2014). Association of untreated metabolic syndrome with moderate to severe periodontitis in Thai population. *Journal of Periodontology*, 85(11), 1502–1514.
- Tonetti, M. S., Greenwell, H., & Kornman, K. S. (2018). Staging and grading of periodontitis: Framework and proposal of a new classification and case definition. *Journal of Periodontology*, 89(Suppl 1), S159–S172.
- Tonetti, M. S., Jepsen, S., Jin, L., & Otomo-Corgel, J. (2017). Impact of the global burden of periodontal diseases on health, nutrition and well-being of mankind: A call for global action. *Journal of Clinical Periodontology*, 44(5), 456–462.
- Wahlin, Å., Papias, A., Jansson, H., & Norderyd, O. (2018). Secular trends over 40 years of periodontal health and disease in individuals aged 20–80 years in Jönköping, Sweden: Repeated cross-sectional studies. *Journal of Clinical Periodontology*, 45(9), 1016–1024.
- Watt, R. G., Daly, B., Allison, P., Macpherson, L. M. D., Venturelli, R., Listl, S., Weyant, R. J., Mathur, M. R., Guarnizo-Herreño, C. C., Celeste, R. K., & Peres, M. A. (2019). Ending the neglect of global oral health: Time for radical action. *The Lancet*, 394(10194), 261–272.
- Wei, Y., Wang, Z., Lei, L., & Chen, L. (2021). Global burden of periodontal disease and its relation with socioeconomic development during 1990–2019. *Journal of Zhejiang University (Medical Science)*, 50(5), 545–552.
- WHO. (2021). *Oral health resolution*. WHO https://apps.who.int/gb/ebwha/pdf_files/EB148/B148_R1-en.pdf.
- Wu, L., Zhang, S., Zhao, L., Ren, Z., & Hu, C. (2022). Global, regional, and national burden of periodontitis from 1990 to 2019: Results from the Global Burden of Disease Study 2019. *Journal of Periodontology*, 93(10), 1445–1454.
- Zaito, T., Kanazawa, T., Shizuma, Y., Oshiro, A., Takehara, S., Ueno, M., & Kawaguchi, Y. (2017). Relationships between occupational and behavioral parameters and oral health status. *Industrial Health*, 55(4), 381–390.
- Zhao, M. J., Qiao, Y. X., Wu, L., Huang, Q., Li, B. H., & Zeng, X. T. (2019). Periodontal disease is associated with increased risk of hypertension: A cross-sectional study. *Frontiers in Physiology*, 10, 440.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Trindade, D., Carvalho, R., Machado, V., Chambrone, L., Mendes, J. J., & Botelho, J. (2023). Prevalence of periodontitis in dentate people between 2011 and 2020: A systematic review and meta-analysis of epidemiological studies. *Journal of Clinical Periodontology*, 1–23. <https://doi.org/10.1111/jcpe.13769>